


Treatment-Naive Genotype 1b Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Patients Genotype 1b Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^a	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks ^b	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for patients who are HIV-uninfected and whose HCV RNA level is <6 million IU/mL	8 weeks ^c	I, B

^a Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

^b For HIV/HCV coinfecting patients, a treatment duration of 12 weeks is recommended.

^c An 8-week regimen can be considered in those with genotype 1b infection and mild fibrosis (see text for details).

Last update: October 24, 2022