

---

## **HCV in Key Populations: Men Who Have Sex With Men**

## Testing

<b>Recommendations for Testing and Prevention of HCV Infection in Men Who Have Sex With Men (MSM)</b>	
RECOMMENDED	RATING
Annual HCV testing is recommended for sexually active HIV-infected adolescent and adult MSM. Depending on the presence of high-risk sexual or drug use practices, more frequent testing may be warranted.	Ila, C
HCV testing at HIV pre-exposure prophylaxis (PrEP) initiation and at least annually thereafter (while on PrEP) is recommended in HIV-uninfected MSM. Depending on sexual or drug use risk practices, more frequent testing may be warranted.	Ila, C
All MSM should be counseled about the risk of sexual HCV transmission with high-risk sexual and drug use practices, and educated about measures to prevent HCV infection or transmission.	Ila, C

## Treatment

<b>Recommendation on Treatment of HCV in Men Who Have Sex With Men (MSM)</b>	
RECOMMENDED	RATING
Antiviral treatment for HCV-infected MSM should be coupled with ongoing counseling about the risk of HCV reinfection, and education about methods to reduce HCV reinfection risk after cure.	I, B

## Testing for HCV Reinfection

<b>Recommendation on Prevention of HCV Reinfection in Men Who Have Sex With Men (MSM)</b>	
RECOMMENDED	RATING
At least annual (and risk-based, if indicated) HCV testing with HCV RNA is recommended for sexually active MSM after successful treatment or spontaneous clearance of HCV infection.	Ila, C

**Last reviewed:** October 24, 2022

