



Treatment-Naive Genotype 3 With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 3 Patients With Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) for patients without baseline NS5A RAS Y93H	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, B
ALTERNATIVE	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) with weight-based ribavirin for patients with baseline NS5A RAS Y93H	12 weeks	IIa, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)/voxilaprevir (100 mg) for patients with baseline NS5A RAS Y93H	12 weeks	IIa, B

^a For [decompensated cirrhosis](#), please refer to the appropriate section.

^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022