


Treatment-Naive Genotype 1a Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 1a Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^a	8 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for patients who are HIV-uninfected and whose HCV RNA level is <6 million IU/mL	8 weeks	I, B
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
ALTERNATIVE	DURATION	RATING
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A


^a Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022

Treatment-Naive Genotype 1a With Compensated Cirrhosis


Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 1a Patients With Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, B

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 1a Patients With Compensated Cirrhosis^a

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
ALTERNATIVE	DURATION	RATING 
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A

^a For [decompensated cirrhosis](#), please refer to the appropriate section.


^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022

Treatment-Naive Genotype 1b Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Patients Genotype 1b Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^a	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks ^b	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for patients who are HIV-uninfected and whose HCV RNA level is <6 million IU/mL	8 weeks ^c	I, B

^a Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

^b For HIV/HCV coinfecting patients, a treatment duration of 12 weeks is recommended.

^c An 8-week regimen can be considered in those with genotype 1b infection and mild fibrosis (see text for details).

Last update: October 24, 2022

Treatment-Naive Genotype 1b With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 1b Patients With Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, B
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A

^a For [decompensated cirrhosis](#), please refer to the appropriate section.

^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022

Treatment-Naive Genotype 2 Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 2 Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^a	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A


^a Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022

Treatment-Naive Genotype 2 With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 2 Patients With Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, B

^a For [decompensated cirrhosis](#), please refer to the appropriate section.


^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022

Treatment-Naive Genotype 3 Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:




Treatment-Naive Genotype 3 Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^a	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A

^a Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

Last update: October 24, 2022

Treatment-Naive Genotype 3 With Compensated Cirrhosis


Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for: Treatment-Naive Genotype 3 Patients With Compensated Cirrhosis^a 		
RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) for patients without baseline NS5A RAS Y93H	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, B
ALTERNATIVE	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) with weight-based ribavirin for patients with baseline NS5A RAS Y93H	12 weeks	IIa, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)/voxilaprevir (100 mg) for patients with baseline NS5A RAS Y93H	12 weeks	IIa, B
<p>^a For decompensated cirrhosis, please refer to the appropriate section.</p> <p>^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.</p>		

Last update: October 24, 2022

Treatment-Naive Genotype 4 Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 4 Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^a	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) ^b	12 weeks	I, A

^a Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.


^b An 8-week regimen can be considered in patients with favorable baseline characteristics (ie, no cirrhosis, HCV RNA <6 million IU/mL, and absence of genotype 4r).

Last update: October 24, 2022

Treatment-Naive Genotype 4 With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 4 Patients With Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks ^c	I, B
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	IIa, B
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	IIa, B

^a For [decompensated cirrhosis](#), please refer to the appropriate section.

^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.


^c For HIV/HCV-coinfected patients, a treatment duration of 12 weeks is recommended.

Last update: October 24, 2022

Treatment-Naive Genotype 5 or 6

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

Treatment-Naive Genotype 5 or 6 Patients With and Without Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, A ^c
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, B
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) ^d	12 weeks	Ila, B

^a For [decompensated cirrhosis](#), please refer to the appropriate section.

^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

^c For compensated cirrhosis, rating is I, B.

^d Not recommended for genotype 6e if subtype is known.

Last update: October 24, 2022